

## Effect of culture temperature on the heterologous expression of *Pleurotus eryngii* versatile peroxidase in *Aspergillus* hosts

G. M. Eibes · T. A. Lú-Chau · F. J. Ruiz-Dueñas ·  
G. Feijoo · M. J. Martínez · A. T. Martínez ·  
J. M. Lema

Received: 24 January 2008 / Accepted: 29 April 2008 / Published online: 15 May 2008  
© Springer-Verlag 2008

**Abstract** Production of recombinant versatile peroxidase in *Aspergillus* hosts was optimized through the modification of temperature during bioreactor cultivations. To further this purpose, the cDNA encoding a versatile peroxidase of *Pleurotus eryngii* was expressed under control of the alcohol dehydrogenase (*alcA*) promoter of *Aspergillus nidulans*. A dependence of recombinant peroxidase production on cultivation temperature was found. Lowering the culture temperature from 28 to 19 °C enhanced the level of active peroxidase 5.8-fold and reduced the effective proteolytic activity twofold. Thus, a maximum peroxidase activity of 466 U L<sup>-1</sup> was reached. The same optimization scheme was applied to a recombinant *Aspergillus niger* that bore the alcohol dehydrogenase regulator (*alcR*), enabling transformation with the peroxidase cDNA under the same *alcA* promoter. However, with this strain, the peroxidase activity was not improved, while the effective proteolytic activity was increased between 3- and 11-fold compared to that obtained with *A. nidulans*.

**Keywords** *Aspergillus* · Proteases · Temperature · Versatile peroxidase

### Introduction

Ligninolytic enzymes have been extensively investigated because of their biotechnological potential for lignin removal in paper pulp manufacturing and in the degradation of aromatic xenobiotics [1–3]. Two ligninolytic peroxidases, lignin peroxidase (LiP) [4, 5] and manganese peroxidase (MnP) [6], were described in *Phanerochaete chrysosporium* and reported in other white-rot fungi from the group of basidiomycetes [7, 8]. A third ligninolytic peroxidase, characterized by its ability to oxidize both Mn<sup>2+</sup> and different aromatic compounds and dyes—including those that are not efficiently oxidized by LiP and MnP, has been reported in *Pleurotus* and *Bjerkandera* species [9–11]. The genes encoding two isoenzymes of this new versatile peroxidase (VP) have been cloned from *Pleurotus eryngii*: *vpl*, expressed in peptone-containing liquid cultures as two allelic variants; and *vpsI*, expressed during lignocellulose solid-state fermentation [12, 13]. *P. eryngii vpl* has been expressed in the model ascomycete *Emericella nidulans* (conidial state *Aspergillus nidulans*) under control of the *alcA* promoter [14]. *A. nidulans* is a genetically well-characterized fungus compared with industrial species such as the deuteromycete-type *Aspergillus niger* and *Aspergillus oryzae*. However, higher levels of recombinant proteins can be obtained from these industrial species [15].

Heterologous expression of active ligninolytic peroxidases from *P. chrysosporium* has been reported in baculovirus systems [16], *Escherichia coli* [17, 18], *Pichia pastoris* [19], *A. oryzae* [20], *A. niger* [21, 22], and *A. nidulans* [23]. However, in no cases have the ligninolytic peroxidase levels been satisfactory, taking into account the amount of enzyme required for its biotechnological application. This low yield has been attributed,

G. M. Eibes · T. A. Lú-Chau (✉) · G. Feijoo · J. M. Lema  
Department of Chemical Engineering, Institute of Technology,  
15706 Santiago de Compostela, Spain  
e-mail: eqtluch@lugo.usc.es

G. M. Eibes  
e-mail: geibes@usc.es

F. J. Ruiz-Dueñas · M. J. Martínez · A. T. Martínez  
Centro de Investigaciones Biológicas, CSIC,  
Ramiro de Maezu 9, 28040 Madrid, Spain

among other reasons, to factors related to the hemoprotein nature of the enzyme, operating at the post-translational level [24]. Some efforts are being directed at better understanding the importance of this aspect of heterologous peroxidase production through the application of genetic engineering techniques [24, 25]. On the other hand, only a few studies have been reported that deal with improving the production of these recombinant enzymes through the optimization of operational conditions.

Both optimization approaches—genetic or operational control improvements—require the identification of the expression bottleneck as a first step. Bioreactor cultivation, in which operational parameters can be easily monitored and controlled, is a very useful tool in identifying such problems. The magnitude of the problem and the feasibility of the optimization of culture conditions may determine whether genetic or process engineering techniques should be used. The optimization of culture conditions seems to be an adequate alternative for improving heterologous peroxidase production, because it is possible to obtain an improvement of the heterologous protein expression by simply optimizing a process variable [26]. However, if a cause–effect relationship could be postulated, this knowledge can be used to improve the heterologous expression by a further modification of the recombinant system. In the present work, the influence of temperature in bioreactor cultivations is considered as a means of improving the production of recombinant *P. eryngii* VP (VP\*) in two *Aspergillus* hosts.

## Materials and methods

### Chemicals

Hemin and threonine (>99%) were obtained from Fluka. Bovine serum albumine (BSA, fraction V) was obtained from Merck. All the components used in growth and induction media were obtained from Sigma-Aldrich.

### Organisms and culture media

The recombinant *A. nidulans* (*biA1*, *methG1*, *argB2*) strain (IJFM A742) harboring the *P. eryngii vpl* cDNA (encoding the allelic variant VPL2) under control of the *alcA* promoter [14] was grown in complete medium, [27] supplemented with D-biotin (10  $\mu\text{g L}^{-1}$ ) and L-methionine (74.5  $\text{mg L}^{-1}$ ). The recombinant *A. niger* (*gpdalCR*, *cspA1*, *leuA1*, *nicA1*, *pyrA6*) strain bearing the *alcA* regulator (*alcR*) of *A. nidulans* was cotransformed according to the method described by Punt and van den Hondel [28] with the same construction used for *A. nidulans* transformation and the vector PAN7-1, which contains the

hygromycin B resistance marker from *E. coli* under the *A. nidulans gpd* gene promoter [29]. The recombinant *A. niger* strain harboring the *vpl* cDNA was grown in glucose–peptone medium [30] supplemented with leucine (250  $\text{mg L}^{-1}$ ) and nicotinic acid (1  $\text{mg L}^{-1}$ ).

Asexual spores (conidia) were removed from Petri dishes by adding 5 mL of sterilized water after a 48 h growth period at 37 °C. Ten milliliters of the spore solution were used to inoculate 1 L of growth medium for experiments with *A. nidulans*. In all other experiments, the initial spore concentration of the growth medium (typically  $1 \times 10^8$  spores  $\text{mL}^{-1}$ ) was fixed by first determining the concentration of the spore solution using a Bürker–Türk counting chamber. Both *Aspergilli* were grown for a period of 24 h at 28 °C and 180 rpm in 1-L Erlenmeyer flasks containing 200 mL of culture medium (growth phase). The washed mycelium was then transferred to the induction medium, with the same composition but containing 0.05% glucose (subinhibitory concentration), 0.1 M threonine (as an inducer), and 0.05% hemin (induction phase). The addition of hemin was required for peroxidase production [14]. Subsequent cultivation of induced cultures was carried out in a bioreactor (as described below).

### Bioreactor cultivation

Bioreactor cultivation was carried out in a BIOSTAT-Q® fermentation system (B. Braun Biotech International, Melsungen, Germany) equipped with four 1-L vessels. pH was kept constant at 6.8 and 5.5 during *A. nidulans* and *A. niger* cultivation, respectively, by the addition of 2 M NaOH or H<sub>2</sub>SO<sub>4</sub> using an automatic control system. The other operational conditions were temperature 19 °C (unless otherwise stated), air flow (2 vvm), and agitation rate (500 rpm, agitated by magnetic stirrers). The initial working volume of all experiments was 0.5 L. When simultaneous experiments were conducted with the same organism, the different vessels were inoculated from a common growth culture. The culture with *A. nidulans* at the optimal temperature was conducted in triplicate in independent fermentations. In all other cases, the cultures were carried out once. Bioreactor cultures of *A. niger* gp20' lacking active copies of the *vpl* gene was inoculated with spore solutions ( $1 \times 10^8$  spores  $\text{mL}^{-1}$ ). All other conditions were similar to those described above.

### Enzymatic activities

VP activity in culture filtrates was estimated from formation of the Mn<sup>3+</sup>-tartrate complex ( $\lambda_{238}$ , 6,500  $\text{M}^{-1} \text{cm}^{-1}$ ) during oxidation of 0.1 mM MnSO<sub>4</sub> in 0.1 M sodium tartrate, pH 5.0, in the presence of 0.1 mM H<sub>2</sub>O<sub>2</sub> [9]. Control measurements without H<sub>2</sub>O<sub>2</sub> were taken. One unit of VP

activity was defined as the amount of enzyme oxidizing 1  $\mu\text{mol}$  of  $\text{Mn}^{2+}$  per minute.


Extracellular protease activity was determined according to the method of van den Hombergh et al. [31]. A 450  $\mu\text{L}$  diluted sample was incubated with 50  $\mu\text{L}$  of 1% (w/v) BSA in 0.1 M sodium acetate buffer (pH 4 and 6.8, for *A. niger* and *A. nidulans* samples, respectively), at 37 °C. After 60 min, reactions were stopped with 500  $\mu\text{L}$  of 10% (w/v) trichloroacetic acid (TCA). Then, the sample and a non-incubated control were kept at 0 °C for 30 min, the precipitated proteins were removed by centrifugation at 6,000 rpm for 5 min, and the optical density of the TCA-soluble fractions was measured at 280 nm. Before calculating protease units, control absorbance was subtracted from sample absorbance. One unit of protease activity was defined as the amount of enzyme causing a change of one absorbance unit per hour.

## Results

### Bioreactor cultivation of recombinant *A. nidulans*

Cultivations of *A. nidulans* have traditionally been performed in the 28–37 °C range [32]. However, it has been shown that, in many cases, the optimum growth conditions are not necessarily the best for the expression of a heterologous enzyme [33]. Previous observations had shown hardly detectable VP activity in shaken cultures of recombinant *A. nidulans* without temperature control, although the temperatures attained (up to 32 °C) did not reduce fungal growth. However, when the temperature was maintained constant at 28 °C, consistent VP activity around 50  $\text{U L}^{-1}$  resulted [26]. In the present work, in order to investigate the effect of temperature on VP\* production, five discontinuous batch bioreactor cultures were conducted at different temperatures (31, 28, 25, 19 and 16 °C), using complete medium [14] (Fig. 1). The highest activity was obtained at 19 °C (466  $\text{U L}^{-1}$ ), while the lowest VP activity was obtained at 31 °C (24  $\text{U L}^{-1}$ ). A correlation between maximal VP activity and the time needed to reach that activity was found (shown in Fig. 1). Thus, the highest and lowest VP activity peaks were obtained after 115 and 14 h, during the fermentations conducted at 19 and 31 °C, respectively.

It can be observed that the temperature effect on VP activity over the experimental range is not symmetrical with respect to that which is optimum for active VP\* production. VP activity slowly decreased when the temperature was increased from 19 to 25 °C, while a sharp activity drop was produced by decreasing temperature from 19 to 16 °C. When independent fermentations at the optimal temperature were repeated in triplicate, a standard deviation of 9.4% of the maximal VP activity was found.




**Fig. 1** Maximal VP activity during bioreactor cultivation of *A. nidulans* at five different temperatures (the growth period required is indicated)

When extracellular protease activity was measured in samples taken from the cultures conducted at the different temperatures, a similar pattern between VP\* production and activity of extracellular proteases was observed (Fig. 2). The maximal level of both MnP and proteolytic activities were reached at the same time, and this time was progressively delayed at lower temperatures.

### Bioreactor cultivation of recombinant *A. niger*

*A. niger* was chosen for the expression of VP\*, since high levels of recombinant proteins can be obtained from this host organism [15]. Since *A. niger* and *A. nidulans* are related species, similar fermentation conditions could be




**Fig. 2** Time course of extracellular VP and protease activities during bioreactor cultivation of *A. nidulans* at 28 °C (filled triangle), 25 °C (filled circle), and 19 °C (filled square)

applied. Moreover, the use of a recombinant *A. niger* strain (bearing *A. nidulans alcR*) enabled VP\* expression under the same *alcA* promoter used in *A. nidulans*. Thus, the next objective was to apply the optimization scheme found for *A. nidulans* to improve the expression of VP\* using *A. niger* as the host organism.

Two simultaneous bioreactor cultures with a recombinant *A. niger* strain harboring the *vpl* cDNA, were conducted at (a) the optimal temperature found for the expression of the *vpl2* peroxidase in *A. nidulans* (19 °C) and (b) the optimal growth temperature of *Aspergillus* (28 °C) (Fig. 3). The highest VP activity was obtained at 19 °C (412 U L<sup>-1</sup>), while at 28 °C only 26% of this activity was found (107 U L<sup>-1</sup>). An inverse correlation between VP activity and extracellular protease level at the two temperatures was observed. This tendency is similar to that found with *A. nidulans* (Fig. 2), but the protease levels were much higher in the recombinant *A. niger* cultures. A third bioreactor was inoculated with *A. niger* spores from a strain lacking VP\* production, which resulted in the absence of extracellular proteases. Since extracellular proteases can strongly affect the production of active VP\*, protease production in both *Aspergillus* hosts was investigated in more detail.

#### Effect of temperature on proteolytic activity of culture broths of *A. niger* and *A. nidulans*

The effect of temperature on extracellular protease activity was investigated using filtered culture broths from




**Fig. 3** Time course of extracellular VP and protease activities during bioreactor cultivation of *A. niger* at 19 °C (filled circle) and 28 °C (filled triangle), as well as *A. niger* gp20' lacking active copies of the *vpl* gene (star)

bioreactor fermentations of *A. nidulans* and *A. niger* (after 7 and 11 days, respectively), each concentrated 30-fold by ultrafiltration (10-kDa cut-off membrane). The protease activity of the concentrated culture broths from *A. nidulans* and *A. niger* fermentations measured at 19 °C were 6,132 and 5,040 U/L, respectively. Extracellular protease activity was analyzed at 37, 31, 28, 25 and 19 °C. In Fig. 4 it can be observed that the activity of the proteolytic enzymes is significantly reduced by lowering the incubation temperature. Thus, *A. nidulans* protease activities at 28 and 19 °C are 38 and 9% of that measured at 37 °C, respectively; while for *A. niger* they are 40 and 19%, respectively. Since routine extracellular protease assays were performed at 37 °C, the actual values of the above results were taken into account to compare the effective proteolytic capacities of *A. nidulans* and *A. niger* cultures grown at different temperatures (Table 1). The lower the temperature of the culture, the lower the production of proteases in both *Aspergillus* strains, this difference being higher for *A. niger* cultures. It is important to highlight that the time necessary to reach the maximal VP activity was double for *A. niger* cultures.

#### Discussion

Operational control during bioreactor cultivation can greatly improve the production of recombinant proteins. In a previous work, we had demonstrated that the production of *P. eryngii* VP\* in *A. nidulans* is considerably reduced as the pH of the medium increases under pH-free conditions [26]. By simply keeping the culture pH constant during bioreactor cultivation, it was possible to improve active VP\* 5.3-fold. The culture temperature used in those experiments was 28 °C, which is considered as the optimal growth temperature of *Aspergillus* sp. In the present study,



**Fig. 4** Estimation of protease activity at different temperatures in samples obtained from bioreactor cultures of *A. niger* and *A. nidulans* (void bars) after 7 and 11 days, respectively (as percentages of the activity measured at 37 °C)

**Table 1** Maximal effective proteolytic activities at 28 and 19 °C during *A. nidulans* and *A. niger* cultures, calculated using the factors obtained from Fig. 4

	<i>A. nidulans</i>		<i>A. niger</i>	
	28	19	28	19
<i>T</i> (°C)	28	19	28	19
Maximal protease activity (U L <sup>-1</sup> )	58	31	633	93
Time (h)	22	114	41	214

The time at which those activities were reached is also presented

we studied the effect of culture temperature on VP\* production using *A. nidulans* and *A. niger* as host organisms. As in previous studies, the *vpl* cDNA encoding a VP isoenzyme expressed in liquid cultures of *P. eryngii* was used for heterologous expression [14].

#### Effect of temperature on VP\* production

Bioreactor cultures at different temperatures demonstrated that a small increase of temperature from 28 °C drastically reduced the production of active VP\*, while active VP\* production was increased 5.8-fold by reducing the temperature from 28 to 19 °C. The effect of culture temperature on heterologous expression has been observed in other recombinant systems [34–36]. Improvement of laccase expression in *Saccharomyces cerevisiae* and of interleukin-3 in *Streptomyces lividans* have previously been obtained by lowering temperature [33, 37]. The opposite effect of temperature has also been reported [38]. However, the present paper is, to the best of our knowledge, the first report on improvement of heterologous expression of a ligninolytic peroxidase by lowering culture temperature.

The effect of temperature on heterologous gene expression has been related to: (a) protein aggregation [39, 40]; (b) incorrect gene translation due to the presence of rarely used codons [41]; and (c) proteolytic degradation [37]. Ligninolytic peroxidases expressed in *Aspergillus* hosts are secreted into the culture medium [14, 20, 22], and no aggregation has been reported. Moreover, codons rarely used by *A. nidulans* (codon usage data from GenBank Release 127.0) were not found in the VP cDNA (GenBank accession No. AF007222). However, as discussed below, a probable relationship between active VP\* production and proteolytic degradation could be established when extracellular proteases were analyzed during recombinant *Aspergillus* cultivation at different temperatures.

#### VP\* production in *Aspergillus niger*

Although VP\* expression was enhanced by decreasing temperature, industrial application of this enzyme (e.g. in paper pulp bleaching or dye decolorization) requires

greater VP\* production. For this reason, the use of other fungal hosts was considered.

Temperature dependence of VP\* production was also observed during *A. niger* bioreactor cultivation. In this case, lowering the cultivation temperature from 28 to 19 °C caused 3.8-fold increase of VP activity. However, when comparing the VP activities produced at 19 °C by the recombinant *A. nidulans* and *A. niger* strains (466 and 412 U L<sup>-1</sup>, respectively), no improvement was achieved by using the second fungal host. The lower VP\* production seems to be due to the high level of extracellular proteases found in *A. niger* cultures. In fact, the effective proteolytic activity in the *A. niger* cultures were 11- and 3-fold those found with *A. nidulans*, when grown at 28 and 19 °C, respectively.

The lack of extracellular proteases during bioreactor cultivation of a recombinant *A. niger* strain that does not produce VP\* suggests that the proteolytic activity is induced by expression of the recombinant peroxidase. It has been shown that proteases are responsible for the degradation of many heterologous proteins [42–44]. Furthermore, in *A. nidulans*, the production of extracellular proteases is subjected to at least four regulatory mechanisms; C, N and S metabolite repression, and pH control. A fifth regulatory mechanism, induction by exogenous protein, is present in *A. niger* [45] and may also operate in *A. nidulans* under certain conditions [46]. The latter mechanism would be responsible for VP\* degradation in both *A. nidulans* and *A. niger* cultures.

The use of protease-deficient hosts has been considered for reducing the problem of proteolytic degradation of heterologous enzymes [22]. Currently, the *vpl* cDNA has been cloned in a protease-deficient *A. niger* strain. However, even using such host organisms, the problem of heterologous protein degradation was not overcome in some cases [44, 47, 48], probably due to the complexity of the *A. niger* protease system [49].

It is noteworthy that related enzymes can be expressed with very different efficiencies in the same or related expression systems. This is the case for basidiomycete peroxidases including: *Coprinus cinereus* peroxidase (CIP), that is expressed with high yields in *A. oryzae* (CIP\* is commercialized by Novozymes, Denmark); *P. chrysosporium* MnP and *Caldariomyces fumago* chloroperoxidase, which can be expressed with low yields in *A. niger* [22, 24, 50] and, in the case of MnP, also in *A. oryzae* [20]; and *P. chrysosporium* LiP, which failed to be expressed in any of the above systems [51]. It has been shown that the *P. eryngii* VP is more related to LiP than MnP in both protein sequence (and molecular structure) and gene regulation [14, 52]. Therefore, the heterologous expression yields reported here, as well as in previous studies [12], are especially relevant, although they should be improved for

biotechnological applications in sectors such as paper pulp bleaching and dye decolorization where high amounts of low price enzyme are required.

## Conclusions

VP\* production in *Aspergillus* hosts can be significantly improved by selecting the optimal culture temperature. Proteolytic degradation of recombinant proteins often represents a limiting factor for increasing heterologous expression. The results obtained suggest that extracellular proteolytic activity involved in VP\* degradation during its expression in *Aspergillus* hosts can be reduced considerably by lowering the cultivation temperature.

**Acknowledgments** The authors thank M. A. Peñalva (CIB, CSIC, Madrid) for providing the *A. nidulans* strain and the plasmid *palcA*, B. Felenbock (University of Orsay, Paris) for the recombinant *A. niger* strain containing the *alcA* regulator, and P. J. Punt (TNO Voeding, Zeist, The Netherlands) for the PN7 vector. This work was partially supported by the EU contract “Fungal metalloenzymes oxidizing aromatic compounds of industrial interest” (QLK3-99-590), the Comunidad de Madrid, and the Spanish Commission of Science and Technology (BIO98-610 and BIO99-908).

## References

- Kirk TK, Farrell RL (1987) Annu Rev Microbiol 41:465–505
- Hammel KE (1992) In: Sigel H, Sigel A (eds) Metal ions in biological systems, vol 28. Degradation of environmental pollutants by microorganisms and their metalloenzymes. Dekker, NY, pp 41–60
- Cai DY, Tien M (1993) J Biotechnol 30:79–90
- Tien M., Kirk TK (1983) Burds Science 221:661–663
- Glenn JK, Morgan MA MB, Kuwahara M, Gold MH (1983) Biochem Biophys Res Commun 114:1077–1083
- Kuwahara M, Glenn JK, Morgan MA, Gold MH (1984) FEBS Lett 169:247–250
- Hatakka A (1994) FEMS Microbiol Rev 13:125–135
- Peláez F, Martínez MJ, Martínez AT (1995) Mycol Res 99:37–42
- Martínez MJ, Ruiz-Dueñas FJ, Guillén F, Martínez AT (1996) Eur J Biochem 237:424–432
- Mester T, Field JA (1998) J Biol Chem 273:15412–15417
- Heinfling A, Ruiz-Dueñas FJ, Martínez MJ, Bergbauer M, Szewzyk U, Martínez AT (1998) FEBS Lett 428:141–146
- Ruiz-Dueñas FJ, Martínez MJ, Martínez AT (1999) Mol Microbiol 31:223–236
- Camarero S, Ruiz-Dueñas FJ, Sarkar S, Martínez MJ, Martínez AT (2000) FEMS Microbiol Lett 191:37–43
- Ruiz-Dueñas FJ, Martínez MJ, Martínez AT (1999) Appl Environ Microbiol 65:4705–4707
- Gouka RJ, Punt PJ, van den Hondel CAMJJ (1997) Appl Microbiol Biotechnol 47:1–11
- Johnson TM, Pease EA, Li JKK, Tien M (1992) Arch Biochem Biophys 296:660–666
- Whitwam R, Tien M (1996) Arch Biochem Biophys 333:439–446
- Doyle WA, Smith AT (1996) Biochem J 315:15–19
- Gu L, Lajoie C, Kelly C (2003) Biotechnol Prog 19:1403–1409
- Stewart P, Whitwam RE, Kersten PJ, Cullen D, Tien M (1996) Appl Environ Microbiol 62:860–864
- Aifa MS, Sayadi S, Gargouri A (1999) Biotechnol Lett 21:849–853
- Conesa A, van den Hondel CAMJJ, Punt PJ (2000) Appl Environ Microbiol 66:3016–3023
- Larrondo LF, Lobos S, Stewart P, Cullen D, Vicuña R (2001) Appl Environ Microbiol 67:2070–2075
- Conesa A, Jeenes D, Archer DB, van den Hondel CAMJJ, Punt PJ (2002) Appl Environ Microbiol 68:846–851
- Elrod SL, Cherry JR, Jones A (2001) Method for increasing hemoprotein production in filamentous fungi. US Patent No. 6,261,827
- Lu-Chau TA, Ruiz-Dueñas FJ, Camarero S, Feijoo G, Martínez MJ, Lema JM, Martínez AT (2004) Bioprocess Biosyst Eng 26:287–293
- Cove DJ (1966) Biochim Biophys Acta 113:51–56
- Punt PJ, van den Hondel CAMJJ (1992) Methods Enzymol 216:447–457
- Punt PJ, Oliver RP, Dingemans MA, Pouwels PH, van den Hondel CAMJJ (1987) Gene 56:117–124
- Kimura Y, Asada Y, Kuwahara M (1990) Appl Microbiol Biotechnol 32:436–442
- van den Hombergh JP, van de Vondervoort PJ, van der Heijden NC, Visser J (1995) Curr Genet 28:299–308
- Calera JA, Lopez-Medrano R, Ovejero MC, Puente P, Leal F (1994) Infect Immun 62:2322–2333
- Cassland P, Jönsson LJ (1999) Appl Microbiol Biotechnol 52:393–400
- Hemila H, Pokkinen M, Palva I (1992) J Biotechnol 26:245–256
- Mongkolsuk S, Loprasert S, Vattanaviboon P, Chanvanichayachai C, Chamnongpol S, Supsamran N (1996) J Bacteriol 178:3578–3584
- Kaufmann H, Mazur X, Fussenegger M, Bailey JE (1999) Biotechnol Bioeng 63:573–582
- Yun SI, Yahya AR, Cossar D, Anderson WA, Scharer JM, Moo-Young M (2001) Biotechnol Lett 23:1903–1905
- Rossini D, Porro D, Brambilla L, Venturini M, Ranzi BM, Vannoni M, Alberghina L (1993) Yeast 9:77–84
- Schein CH, Noteborn MHM (1988) Biotechnology 6:291–294
- Vasina JA, Baneyx F (1997) Protein Expr Purif 9:211–218
- Boule JB, Papanicolaou C, Rougeon (2000) Enhanced expression of heterologous proteins in recombinant bacteria through reduced growth temperature and co-expression of rare tRNA's. European Patent EP1010763
- Roberts IN, Jeenes DJ, MacKenzie DA, Wilkinson AP, Sumner IG, Archer DB (1992) Gene 122:155–161
- Broekhuijsen MP, Mattern IE, Contreras R, Kinghorn JR, van den Hondel CA (1993) J Biotechnol 31:135–145
- Krasevec N, van de Hondel CA, Komel R (2000) Pflugers Arch 440(Suppl 5):R83–R85
- Jarai G, Buxton F (1994) Curr Genet 26:238–244
- Katz ME, Masoumi A, Burrows SR, Shirliff CG, Cheetham BF (2000) Genetics 156:1559–1571
- Mattern IE, van Noort JM, van den Berg P, Archer DB, Roberts IN, van den Hondel CA (1992) Mol Gen Genet 234:332–336
- Wiebe MG, Karandikar A, Robson GD, Trinci AP, Candia JL, Trappe S, Wallis G, Rinas U, Derckx PM, Madrid SM, Sisniega H, Faus I, Montijn R, van den Hondel CA, Punt PJ (2001) Biotechnol Bioeng 76:164–174
- van den Hombergh JP, Fraissinet-Tachet L, van de Vondervoort PJ, Visser J (1997) Curr Genet 32:73–81
- Conesa A, van De Velde F, van Rantwijk F, Sheldon RA, van Den Hondel CA, Punt PJ (2001) J Biol Chem 276:17635–17640
- Conesa A, Punt PJ, van den Hondel CAMJJ (2002) J Biotechnol 93:143–158
- Ruiz-Dueñas FJ, Guillén F, Camarero S, Pérez-Boada M, Martínez MJ, Martínez AT (1999) Appl Environ Microbiol 65:4458–4463